Lal, S. (2025). Analyzing the Impact of Social Isolation on Aging: Health, Mental State and Cognitive Functions. *Critical Debates in Humanities, Science and Global Justice*, *5*(2), 26–37.

RESEARCH ARTICLES

Analyzing the Impact of Social Isolation on Aging: Health, Mental State and Cognitive Functions

Sakhi Lal[a]

Keywords: mental health, aging, well-being

Critical Debates in Humanities, Science and Global Justice

Vol. 5, Issue 2, 2025

Social isolation in older adults has become a significant concern that bears down on physical health, mental health, and cognition. This qualitative research explores the complex consequences of social isolation on older adults. A systematic approach is adopted through semi-structured interviews with fifty individuals in the 65 to 80 age group and a comprehensive review of relevant literature. The results mirror that social isolation is associated with increased susceptibility to chronic diseases, mental illness, and cognitive impairment. The primary themes outlined are the negative impact of reduced social interaction on physical well-being, the worsening of psychological illness, including depression and anxiety, and the speeding up of cognitive decline. The research emphasizes the need for social engagement to counteract these negative impacts and underscores the necessity of targeted interventions in strengthening social support systems among the elderly. From the physical, emotional, and cognitive aspects of social isolation, this study offers insightful contributions to building holistic solutions to facilitate healthy aging and improve well-being in older adults.

Introduction

The global population is experiencing a significant demographic shift towards increased aging, bringing various issues older adults face into the spotlight. Among these, loneliness and social isolation have emerged as prominent concerns. Improved medical care and lifestyle changes have increased survival rates, resulting in a growing number of individuals undergoing prolonged old age. However, this extended lifespan often changes social relations, leading to reduced social interaction due to retirement, the death of a spouse or peers, mobility issues, and geographical distance from relatives. Social isolation, defined as the absence of social contact, interaction, and fulfilling relationships with family, friends, or society in general, is an objective condition that tends to result in loneliness, an emotional state of feeling disconnected or isolated. Interestingly, loneliness can occur even among people, as it is a matter of individual perception and the quality of relationships rather than the number of social contacts.

a Sakhi Lal is a student at Lafayette High School in Missouri.

Despite recognizing social isolation as a significant issue, its precise effects on aging remain poorly understood. Existing research has highlighted various negative impacts, including increased risks for cardiovascular diseases, obesity, and mortality, as well as mental health disorders such as depression, anxiety, and stress. However, there is a need for a more comprehensive understanding of how social isolation affects physical health, mental well-being, and cognitive functioning among the elderly population. This study aims to fill this gap by investigating the multifaceted effects of social isolation on aging. Combining semi-structured interviews with elderly participants and an extensive review of existing literature, this research provides a holistic view of the impact of social isolation. The findings will contribute to developing targeted interventions to enhance social support networks and promote healthy aging.

Background of Study

Social isolation significantly impacts the progression of aging, affecting physical health, mental well-being, and cognitive functioning. It contributes to cardiovascular diseases, obesity, and mortality, and can be mitigated through interventions targeting social connections (U.S. Department of Health and Human Services, 2023). Mental health disorders like depression, anxiety, and stress are also affected, and coping mechanisms or resilience factors may buffer these effects (Hwang et al., 2022). Cognitive function is affected, and social engagement can enhance cognitive functioning. Social isolation also affects healthcare utilization, with economic implications. Social isolation has considerable effects on physical and psychological health, leading to higher medical use, increased cost to healthcare providers, and financial implications for insurance companies. Higher levels of social isolation were significantly associated with increased odds of nursing home placement (odds ratio 2.01) and skilled nursing facility stays (odds ratio 1.16) over 2 years (Pomeroy et al., 2023). Loneliness and perceived social support mediate the relationship, and strategies to enhance social support networks can be implemented. Loneliness, measured by a 3-item loneliness scale, predicted slightly lower Medicare spending of $768 per beneficiary annually, with less inpatient and outpatient care utilization (Shaw et al., 2017). Effective intervention strategies, such as technology and virtual social connections, can help promote healthy aging and well-being in older adults (Jimenez et al., 2023). Health-related information gathered from the sampled populations indicated a high prevalence of chronic medical ailments in the 65- to 80-year-old population. Some of the ailments noted were hypertension, joint pain and back pain, diabetes, arthritis, limited mobility, heart disease, COPD, osteoporosis, high cholesterol, Parkinson's disease, and chronic fatigue syndrome. The severity of these ailments was relatively diverse, with some needing help in daily activities, while others had relatively active lifestyles. Milder medical conditions, including controlled hypertension, occasional headache, and mild asthma, were also seen and generally treated with pharmacologic treatment. How does social isolation contribute to physical

and mental health development and progression? If it does, what role can social engagement play in maintaining cognitive health and preventing cognitive decline in aging populations?

Research Design

This study employed a combination of participant interviews and online research to gather comprehensive data. Initially, a survey was conducted via SurveyMonkey.com to identify potential participants. The survey respondents selected a diverse group of individuals aged 65 to 80 for in-person interviews. Each interview lasted approximately 30 to 45 minutes and was audio-recorded and transcribed for analysis. Concurrently, an extensive online review of academic journals, books, reputable websites, and databases was conducted to provide contextual background and support the interview findings. Thematic analysis was applied to both the interview transcripts and the literature to identify recurring patterns and themes, ensuring a thorough understanding of the research topic. Researchers such as Prita Bhandari, Saul McLeod, and H.M. Levitt have highlighted the success of qualitative research methods. Bhandari (2020) suggests the importance of understanding complicated social phenomena. McLeod (2023) explains the dependability and trustworthiness of qualitative research approaches. Levitt (2020) speaks about the close attention to the subject, suggesting the importance of qualitative research in capturing the richness of human experience.

Literature Review

Social isolation has been associated with a variety of negative physical health consequences. The National Institute on Aging (NIA) reports that social isolation and loneliness are linked with a higher risk of chronic diseases such as hypertension, cardiovascular disease, obesity, and weakened immune function (2019). All these conditions are exacerbated by inflammation, a major driver of chronic disease development. The Centers for Disease Control and Prevention (CDC) emphasizes the important role of social relationships in maintaining physical health (2023).

The NIA notes that loneliness and social isolation can lead to anxiety, depression, and cognitive decline (2019). However, the relationship between mental health and social isolation is complex and multidimensional. According to some studies, coping and resilience factors may function as buffers to these consequences. Hwang et al. demonstrated that social isolation and anxiety in individuals with cognitive impairment are indeed related, but that the presence of social support can buffer these adverse effects. This indicates a requirement for more intricate explorations of the variables that determine mental health results in socially isolated individuals.

A review published in Frontiers in Aging Neuroscience discovered that social isolation and loneliness are associated with poor cognitive health, including declines in memory, attention, and overall cognitive functioning (Cardona & Andrés, 2023). The University of Chicago study emphasizes that the absence of human interaction can lead to significant cognitive losses

(Roy et al., 2023). However, there is conflicting evidence regarding the extent to which social isolation directly causes cognitive decline. Certain studies indicate that loneliness, rather than social isolation, has a more significant correlation with cognitive impairment. For example, Freak-Poli et al. (2022) established that loneliness, rather than social support, is what correlates with cognitive decline and dementia. This indicates that more research needs to be done to determine the unique effects of social isolation and loneliness on cognition.

Social isolation also affects healthcare utilization, with significant economic implications. Higher levels of social isolation are associated with increased odds of nursing home placement and skilled nursing facility stays. Shaw et al. (2017) found that objective social isolation increases Medicare spending, while loneliness does not. This suggests that social isolation leads to higher healthcare costs due to increased medical use. Effective intervention strategies are crucial for mitigating the adverse effects of social isolation. Technology and virtual social connections have been proposed as potential solutions. Jimenez et al. (2023) highlight the importance of personalizing digital health systems to meet the needs of different older populations.

Methodology

Data were collected through two sources: an online survey of literature and in-person interviews. The literature survey included academic journals, reports, and publications to develop a theoretical foundation of social isolation among older adults. Recruitment started with a survey posted through friends, family, and senior residential complexes on SurveyMonkey.com and completed by 450 individuals. A diverse group (Table 1) of people aged 65- 82 with varying living conditions and demographics was selected based on the survey results. Two groups were identified: high and low activity rates. This sample allowed for a valid comparison between socially engaged and withdrawn participants in varying living situations.

I conducted semi-structured interviews with all the participants through a guideline of ten open-ended questions about daily routines and life so that they could provide their experiences at liberty. Face-to-face interviews were conducted over 30-45 minutes, averaging 35 minutes, at senior centers or during home visits. The participants provided informed consent, and all sessions were recorded for precise data analysis.

Technology had a significant role in data processing. Google Speech was used to transcribe audio into a Word document. The text was color-coded to differentiate between my voice and the participant's voice. Data coding used a free trial of MAXQDA, which assisted in categorizing and sentiment analysis (positive, negative, or neutral), through which themes and patterns were determined. All the respondents were asked the same pre-determined questions to maintain consistency. They were paraphrased, coded, and compared for similarity and contrast. The findings were sequenced systematically within the research design to highlight vital themes. Social

Table 1.

Living Status	No	Marital Status	No	Ethnicity	No
Independent Living	25	Married	8	African American	13
Senior Center Living	25	Divorced	7	Hispanic	9
		Single	5	Caucasian	23
		Widow/Widower	30	Asian	5

isolation was quantified using three measures: living alone, interaction frequency, and activity participation. Interview data analysis included pattern and theme identification. Seven overarching themes emerged: Social Connections, Health Limitations, Routine Control, Emotional Well-being, Living Environment Satisfaction, Need for Personalized Care, and Coping with Loss. Thirteen individual patterns emerged within the themes. Interviews and secondary data provided insight into the reasons for social isolation among older people. The following section will present the findings and implications of this examination.

Results

This qualitative research suggests a positive relationship between social interaction and the general health of older people. The analysis contrasts two different cohorts: one characterized by high levels of social isolation, with more incidences of loneliness, depression, anxiety, and a variety of chronic physical health issues; the other characterized by regular social activity in book clubs, community theater, and volunteering. Individuals in the second group had considerably more emotional well-being, such as variables of happiness, contentment, and feeling fulfilled. The model suggests that social engagement shapes older adults' emotional and mental well-being. Higher degrees of social interaction appear to be related to greater psychological outcomes and can be used to model the better management of chronic health. Preliminary results also suggest that those with higher levels of social engagement are more likely to use coping techniques, such as mindfulness and meditation; however, further empirical evidence is needed to establish this relationship. The research recognizes various coping responses among older people, which can be classified into social and individual coping responses. Individual coping responses consisted of creative activities (painting, writing, knitting), mental stimulation (puzzles, reading), passive entertainment (television and radio), and participation in online support groups. Furthermore, some respondents engaged in socially integrative activities, including club membership, group events, and volunteering, thus highlighting the vital protective function of social contact in preserving mental health. Interestingly, several participants who lived alone but described a strong engagement in creative or spiritual activities—such as poetry writing, religious study, gardening, or crafting—reported a surprisingly low level of emotional distress. These individuals frequently described their daily routines with words like "fulfilling," "purposeful," and "joyful." This suggests that a

sense of internal purpose and meaningful engagement may act as a protective buffer, even in the absence of regular social interaction. Such findings introduce the possibility that it is not solely social isolation, but rather the absence of purpose-driven activity, that exacerbates vulnerability to emotional and cognitive decline in older adults. These patterns merit further empirical inquiry.

A comparative examination of coping strategies illustrates substantial disparity between the two samples. Participants who reported high loneliness, depression, and anxiety levels mainly depended on individual activities (such as reading, puzzles, and creative interests) and online support networks (like internet forums and electronic book clubs) as their primary coping strategies. Conversely, individuals with good social networks reported greater emotional well-being and employed socially oriented strategies, including regular exercise (e.g., yoga and swimming), group activity involvement (e.g., volunteers and club membership), and mindfulness. Both groups utilized electronic communication tools like social media and video conferencing, although these tools had different purposes based on everyone's level of social engagement.

The hypothesized comparative model proposes the possibility of a connection between active social involvement and greater mental well-being among older individuals. However, final causation is not warranted based on present data. The difference in emotional experiences and coping mechanisms underlines the complexity of psychological resilience and adaptation in the context of old age. The research also gives a clear overview of the various ways in which older people engage socially. In-person activities involved engaging in special-interest groups (e.g., gardening, chess, photography, bridge, travel, history, and dance) and attending community-based religious services and volunteer activities (e.g., food banks, shelters, and senior centers). Besides, electronic engagement was also significant and involved engaging in online forums, virtual clubs, writing groups, language exchange programs, and poetry readings. Frequent communication with family and friends via telephone and video calls was commonly reported as a means of social contact. Moreover, most participants identified leisure pursuits, such as reading, crafting, gardening, listening to audiobooks and music, and birdwatching, as significant and calming activities. The results indicate that social engagement in old age is multifaceted and multidimensional, consisting of a combination of group activities that are organized and autonomous pursuits, supported by both physical and virtual spaces. To consolidate these findings, follow-up studies must include other data sources and analytical approaches. The research implies a link between social isolation and poor physical health. Individuals who indicated restricted social activity were more prone to chronic health conditions like hypertension, joint and back pain, diabetes, arthritis, and heart disease. Respondents with active social lives and sound interpersonal support systems were likelier to report better physical and mental health outcomes. Although the results show a significant link

between social connectedness and physical health, the research does not make a causal link. Additional longitudinal research is needed to ascertain whether social isolation actively develops or enhances certain diseases.

One of the key limitations of the research is the small sample size (n=50), which limits the potential for undertaking detailed comparative analysis between demographic subgroups. Notwithstanding this limitation, the prevalence of loneliness, depression, and anxiety was high, and a large number of participants were affected by one or more of these mental health issues simultaneously. The findings imply a potential link between social resource availability and coping strategy effectiveness. Future studies will be positioned to investigate this connection more completely, assessing the impact of socioeconomic status, geographical location, and community resource accessibility on mental health outcomes and coping efficacy. Moreover, studies drawing on a larger, more demographically representative sample would increase the generalizability of results. Although the dataset may be revealing, its limited size constrains broader conclusions. Confirmatory documentation and additional data gathering are necessary to validate observed patterns. Future studies could benefit from including alternative methodologies, additional age cohorts, or geographic diversity to give a broader snapshot of health and well-being among older ages. This research identifies the importance of social interaction for older people's mental and physical well-being. Although the results provide useful preliminary information, the small sample size and qualitative nature highlight the requirement for larger and more systematic research to confirm and build on these results.

Discussion

Social interactions deeply influence the mental and physical well-being of older adults. The observations made in this research reflect a clear difference between socially engaged participants and socially isolated participants: socially active participants are satisfied and happy, whereas socially isolated participants are lonely, depressed, suffer from anxiety, and suffer from chronic physical illness. These findings confirm previous research showing that social connectedness is a robust buffer for mental health during old age. The model emerging from this research implies that individuals with greater social engagement are more likely to practice adaptive coping strategies, such as mindfulness, volunteering, and physical activity that enhance resilience to emotional upset and chronic illness. Older people utilize two broadly categorized strategies of coping with isolation: social and individual. Socially active older people get involved in community-based activities (e.g., book clubs, church services, volunteering) and use online media to increase their social network. Conversely, isolated older people engage in individual activities and online support groups, which, while helpful, may not have the emotional intensity of face-to-face contact. A compelling avenue for deeper inquiry lies in questioning whether the adverse outcomes observed in socially isolated individuals stem from the isolation itself or, more precisely, from

a lack of meaningful purpose and engagement in their daily lives. Several participants who had limited social contact yet maintained active, internally motivated routines—such as journaling, painting, mentoring online, or practicing yoga, demonstrated high emotional resilience and mental clarity. These observations raise the possibility that purposeful solitude can offer protective effects like social connection. This aligns with psychological theories emphasizing meaning-making as a driver of well-being and suggests that interventions should not only promote external interaction but also foster individual pursuits that cultivate a sense of purpose, commitment, and inner fulfillment.

Impact of Social Isolation on Physical Well-Being

A startling trend that the research unfolds is the proposed connection between physical decline and social withdrawal. Participants with minimal social contact had higher rates of chronic illnesses like hypertension, arthritis, and heart disease. Though the findings indicate the possibility of a connection between isolation and declining physical well-being, the cross-sectional design of the research restricts definitive causal conclusions. Longitudinal studies are required to determine whether social engagement results in better health or whether healthier individuals are more socially engaged. A key mechanism by which social isolation influences physical health is via chronic stress. Chronic psychological distress activates the body's stress response, prompting increased cortisol release, with potential systemic inflammation consequently. Inflammation has been implicated in cardiovascular disease, hypertension, insulin resistance, and weakened immune function, rendering socially isolated older adults at risk for infection, delayed wound healing, and unfavorable health outcomes (Doane & Adam, 2010). Isolation is associated with physical inactivity. Socially isolated older adults are less likely to be regular physical exercisers (Irvine et al., 2022), which accelerates the rate of loss of muscle mass, reduces mobility, and increases the risk of falls and frailty (Milanović et al., 2013). In addition, social isolation can be responsible for poor adherence to medication regimes, worsening chronic illness over time (Bartley et al., 2024). Social isolation can result in detrimental behaviors, including poor diet, lack of exercise, and failure to obtain medical attention factors all leading to the progression of disease. Thus, the alleviation of social isolation is essential for both the physical and mental well-being of the elderly. Community-based interventions, including organized exercise programs and social support networks, can mitigate the adverse consequences of isolation and promote longer, healthier living (Irvine et al., 2022).

Impact of Social Isolation on Physical Well-Being

Social isolation in mental health is a strong predictor of long-term emotional difficulties, such as depression, anxiety, and despair. Older adults who experience social isolation are more likely to develop major mental illnesses, such as major depressive disorder and generalized anxiety disorder

(New Jersey Department of Human Services, 2023), both of which decrease quality of life and lead to poor physical health and poor cognitive functioning. One of the primary reasons for poor mental health is the lack of emotional support and quality interpersonal relationships (Freak-Poli et al., 2022). Regular social interaction is a cushion against stress, providing company and a feeling of belonging. In its absence, psychological pain may be enhanced, motivation towards self-care might be lost, and overall mental health could be compromised.

Cognitive Consequences of Social Isolation

Cognitive health is also significantly affected by social isolation. Older people who lack frequent social contact are at greater risk of cognitive impairment and neurodegenerative disorders such as Alzheimer's and dementia (Shafighi et al., 2023). Social activity stimulates the brain, maintaining memory, communication, and problem-solving skills. Insufficient mental stimulation can accelerate cognitive impairment and dependence. Unhealthy lifestyle habits that frequently accompany social isolation also increase cognitive risk. Unhealthy diets, lack of exercise, and postponed medical treatment are prevalent among socially isolated individuals. In addition, having fewer social contacts can result in early cognitive decline going undetected, with interventions postponed.

Social isolation decreases stimulation through cognitively demanding activities, like conversation, decision-making, and problem-solving, which decline brain function under the precept of "use it or lose it" (Shen et al., 2022). Furthermore, depression, which is strongly associated with social isolation, can influence this process by modifying neurotransmitter equilibrium and impacting brain morphology, especially in areas crucial in memory and executive functions (Bartley et al., 2023). Solution to these cognitive difficulties necessitates purposeful attempts at facilitating mental stimulation, providing emotional support, and enhancing social engagement of older individuals. Promoting involvement in community and intergenerational activities and using technology for remote contact and cognitive training may preserve cognitive well-being (Bartley et. al.).

Social relationships are most valuable in enhancing quality of life, combating feelings of loneliness, and postponing age-related challenges. Study participants often discussed emotional upset brought about by rare visits from family members or the death of close relatives. Conversely, individuals with strong and good social connections had considerably more positive emotional outlooks. Most effectively to address the consequences of social isolation, interventions should be multimodal. This includes more access to mental health care, greater availability of community-based opportunities for activity engagement, and incentives for digital connectivity solutions (Jimenez et al., 2023). Senior clubs, support groups, and volunteerism can help create meaningful relationships and reduce loneliness (Irvine et al., 2022). One-on-one mental health counseling and screening services can also identify and treat early symptoms of distress, before severe impairment.

Overall, while in its early stages, this study contributes to the accumulating evidence that supports the critical role of social engagement in fostering emotional resilience, cognitive health, and physical well-being in older adults. Future studies should emphasize the necessity for larger and more varied samples and longitudinal mixed methods designs to examine the evolving relationship between social determinants and health over time. Variations of social relationships, such as face-to-face relationships, computer-based communication, and home-based relationships, has been established as one of the core determinants of health outcomes. With growing digital competency among older adults, more research must be conducted on the impact of different social interaction modalities on emotional and physical health.

Conclusions

Social engagement is a central element in physical health promotion and enhancement, as it inspires participation in group exercise sessions, synchronized regimens, and walking in groups, which generate encouragement and enjoyment. Aside from its physiological benefits, socialization is a very powerful emotional support system that helps older adults cope with feelings of depression, loss of independence, and the challenges of aging by facilitating a strong sense of community and belongingness. Engagement in recreational activities, religious practices, and significant interaction with family members also increases mental strength and develops a deep sense of joy and general life satisfaction.

To promote the well-being of older persons, assisted living communities should provide a range of social activities, encourage family involvement, engage in flexible communication approaches, and incorporate intergenerational programs. At the same time, activities based on individual interests can play a key role in stimulating engagement. Improving the well-being of aging populations, therefore, may require a two-pronged approach: the development of social connections and the facilitation of purpose-driven solitary engagement. Recent studies suggest that a strong sense of purpose, irrespective of the level of social engagement, can independently protect emotional and cognitive well-being (Frankl, 2006; Hill & Turiano, 2014). Prioritizing both social and purpose-driven individual activities in eldercare has the potential to lead to improved quality of life and a more meaningful experience of aging.

Published: August 10, 2025 EDT.

This is an open-access article distributed under the terms of the Creative Commons Attribution 4.0 International License (CCBY-4.0). View this license's legal deed at http://creativecommons.org/licenses/by/4.0 and legal code at http://creativecommons.org/licenses/by/4.0/legalcode for more information.

References

Bartley, M. M., St. Sauver, J. L., Baer-Benson, H., Schroeder, D. R., Khera, N., Fortune, E., & Griffin, J. M. (2023). Exploring social determinants of health and physical activity levels in older adults living with mild cognitive impairment and dementia in the Upper Midwest of the United States. *Preventive Medicine*, *177*, 107773. https://doi.org/10.1016/j.ypmed.2023.107773

Bartley, M. M., St. Sauver, J. L., Schroeder, D. R., Khera, N., & Griffin, J. M. (2024). Social isolation and healthcare utilization in older adults living with dementia and mild cognitive impairment in the United States. *Innovation in Aging*, *8*(10), igae081. https://doi.org/10.1093/geroni/igae081

Bhandari, P. (2020, June 19). *What is qualitative research?* Scribbr. https://www.scribbr.com/methodology/qualitative-research/

Cardona, M., & Andrés, P. (2023). Are social isolation and loneliness associated with cognitive decline in ageing? *Frontiers in Aging Neuroscience*, *15*. https://doi.org/10.3389/fnagi.2023.1075563

Doane, L. D., & Adam, E. K. (2010). Loneliness and cortisol: Momentary, day-to-day, and trait associations. *Health Psychology*, *29*(4), 364–373. https://doi.org/10.1037/a0018640

Frankl, V. E. (2006). *Man's Search for Meaning* (Beacon Press ed.). Beacon Press.

Freak-Poli, R., Wagemaker, N., Wang, R., Lysen, T. S., Ikram, M. A., Vernooij, M. W., ... Tiemeier, H. (2022). Loneliness, not social support, is associated with cognitive decline and dementia across two longitudinal population-based cohorts. *Journal of Alzheimer's Disease*, *85*(3), 295–308. https://doi.org/10.3233/JAD-210330

Hill, P. L., & Turiano, N. A. (2014). Purpose in life as a predictor of mortality across adulthood. *Psychological Science*, *25*(7), 1482–1486. https://doi.org/10.1177/0956797614531799

Hwang, Y., Massimo, L., Aryal, S., & Hodgson, N. A. (2022). The relationship between social isolation and anxiety in people with cognitive impairment in the United States. *International Journal of Geriatric Psychiatry*, *37*(2). https://doi.org/10.1002/gps.5679

Irvine, K. N., Fisher, D., Marselle, M. R., Currie, M., Colley, K., & Warber, S. L. (2022). Social isolation in older adults: A qualitative study on the social dimensions of group outdoor health walks. *International Journal of Environmental Research and Public Health*, *19*(9), 5353. https://doi.org/10.3390/ijerph19095353

Jimenez, J., del Rio, A., Berman, A. N., & Grande, M. (2023). Personalizing digital health: Adapting health technology systems to meet the needs of different older populations. *Healthcare*, *11*, 2140. https://doi.org/10.3390/healthcare11152140

Levitt, H. M. (2020). *Reporting qualitative research in psychology: How to meet APA style journal article reporting standards* (Rev. ed.). American Psychological Association. https://doi.org/10.1037/0000179-000

McLeod, S. (2023, December 18). *Qualitative vs quantitative research methods & data analysis*. Simply Psychology. https://www.simplypsychology.org/qualitative-quantitative.html

Milanović, Z., Pantelić, S., Trajković, N., Sporiš, G., Kostić, R., & James, N. (2013). Age-related decrease in physical activity and functional fitness among elderly men and women. *Clinical Interventions in Aging*, *8*, 549–556. https://doi.org/10.2147/CIA.S44112

New Jersey Department of Human Services, Office of Research & Evaluation. (2023). *Social isolation study*.

Pomeroy, M. L., Cudjoe, T. K. M., Cuellar, A. E., Ihara, E. S., Ornstein, K. A., Bollens-Lund, E., ... Gimm, G. W. (2023). Association of social isolation with hospitalization and nursing home entry among community-dwelling older adults. *JAMA Internal Medicine*. https://doi.org/10.1001/jamainternmed.2023.3064

Roy, K., Smilowitz, S., Bhatt, S., & Conroy, M. L. (2023). Impact of social isolation and loneliness in older adults: Current understanding and future directions. *Current Geriatrics Reports*, *12*, 138–148. https://doi.org/10.1007/s13670-023-00393-5

Shafighi, K., Villeneuve, S., Rosa Neto, P., Badhwar, A., Poirier, J., ... Sharma, V. (2023). Social isolation is linked to classical risk factors of Alzheimer's disease-related dementias. *PLoS ONE*, *18*(2), e0280471. https://doi.org/10.1371/journal.pone.0280471

Shaw, J. G., Farid, M., Noel-Miller, C., Joseph, N., Houser, A., Asch, S. M., ... Flowers, L. (2017). Social isolation and Medicare spending: Among older adults, objective social isolation increases expenditures while loneliness does not. *Journal of Aging and Health*, *30*(10), 1497–1513. https://doi.org/10.1177/0898264317703559

Shen, C., Rolls, E. T., Cheng, W., Kang, J., Dong, G., Xie, C., ... Feng, J. (2022). Associations of social isolation and loneliness with later dementia. *Neurology*, *99*(2), e164–e175. https://doi.org/10.1212/WNL.0000000000200583

U.S. Department of Health and Human Services. (2023). *Our epidemic of loneliness and isolation: The U.S. Surgeon General's advisory on the healing effects of social connection and community*. https://www.hhs.gov/sites/default/files/surgeon-general-social-connection-advisory.pdf